



*National Aeronautics and Space  
Administration Goddard Earth Science Data  
Information and Services Center (GES DISC)*

# README Document for Global Precipitation Climatology Project Version 3.0 Precipitation Data

---

**Last Revised: March 26, 2021**

Goddard Earth Sciences Data and Information Services Center (GES DISC)

<https://disc.gsfc.nasa.gov>

NASA Goddard Space Flight Center

Code 610.2

Greenbelt, MD 20771 USA

**Prepared By:**

***GES DISC MEaSURES Team***

*George J. Huffman*

---

Name  
GES DISC  
GSFC Code 610.2

---

Name  
Code 612  
NASA Goddard Space Flight Center

March 14, 2019

---

Date

**Reviewed By:**

*Zhong Liu*

March 26, 2021

---

Reviewer Name  
GES DISC  
GSFC Code 610.2

---

Date

**Goddard Space Flight Center  
Greenbelt, Maryland**

# Revision History

---

<i>Revision Date</i>	<i>Changes</i>	<i>Author</i>
March 14, 2019	Original	George Huffman
March 26, 2021	GES DISC Help Desk new email	Zhong Liu

# Table of Contents

---

## Contents

1.0 Introduction.....	6
1.1 Dataset/Mission Instrument Description .....	6
1.1.1 GPCPV3.0.....	7
1.2 Algorithm Background .....	7
1.3 Data Disclaimer .....	8
1.3.1 Data Citation .....	9
1.3.2 Contact Information .....	9
1.4 What's New? .....	9
2.0 Data Organization .....	9
2.1 File Naming Convention .....	10
2.2 File Format and Structure.....	10
2.3 Key Science Data Fields .....	10
3.0 Data Contents .....	10
3.1 Dimensions .....	10
3.2 Global Attributes.....	10
3.3 Products/Parameters .....	13
4.0 Options for Reading the Data .....	13
4.1 Command Line Utilities .....	13
4.2 Tools/Programming .....	14
5.0 Data Services .....	14
6.0 More Information .....	15
7.0 Acknowledgements .....	15
References .....	15

## Tables

Table 1: Global metadata attributes associated with each SDS. ....	11-12
--	-------

Table 2: Key Metadata Items. .... 12

Table 3: Data fields in the GPCP V3.0 Monthly dataset. The latitude/longitude use the WGS84 reference datum. .... 13

# 1.0 Introduction

---

Precipitation observations are critical to research and many applications, including drought monitoring, flash floods, crop forecasting, disease prediction, and ocean salinity studies. Rain gauges are the primary source of direct precipitation observations. Unfortunately, rain gauges are point measurements and much of the globe is sparsely covered, especially in underdeveloped countries and areas of low population density. Furthermore, with the exception of a few buoy arrays, there are no precipitation gauge observations over the open ocean. Satellites seek to mitigate the limitations of rain gauge observations by estimating precipitation over land and ocean for most, or all, of the entire globe. When converted to gridded precipitation estimates, the satellite observations facilitate a multitude of studies, including those on the larger space-time scales that gauge analyses typically cannot provide. To augment the satellite-based precipitation estimates, uniformly processed gauge analyses are incorporated to improve the land-based estimates.

The Global Precipitation Climatology Project (GPCP) is a community-based activity of the Global Water and Energy Exchange (GEWEX) project in the World Climate Research Programme (WCRP), focused on creating a global, long-term homogeneous record of gridded precipitation estimates and ancillary information for use in climate studies and other applications. GPCP Version 3 is the successor to the highly successful GPCPV2 data set. The current GPCPV3.0 products are at the monthly resolution with the daily and 3-hourly products to be developed next.

This first release, labeled V3.0, is considered to be a beta release, meaning it is useful, but has known limitations. Specifically, the TOVS/AIRS record is not as homogeneous as we expect for a CDR, and the IR input has some noticeable seams. Users should account for these limitations in the use of the V3.0 data. The team continues to work toward improving these issues in a future release.

## 1.1 Dataset/Mission Instrument Description

The GPCPV3.0 data set provides a gridded (Level 3) homogeneously processed record of global precipitation estimates at  $0.5^\circ$  spatial and monthly temporal resolution. The current data span is (1983–2016) with the potential to extend this record in the future. The current monthly product provides the following data fields:

- (1) merged satellite-gauge precipitation estimate,
- (2) merged satellite-gauge precipitation random error estimate,
- (3) satellite-only precipitation estimate,

- (4) satellite source field,
- (5) gauge analysis precipitation, and
- (6) probability of liquid-phase precipitation.

### 1.1.1 GPCPV3

The GPCPV3.0 estimates rely on a relatively homogeneous record of satellite precipitation estimates and gauge analyses as input. Passive microwave (PMW) radiometer estimates are computed from the Defense Meteorological Satellite Program (DMSP) Special Sensor Microwave/Imager (SSM/I) that flew on the F11 and F13 spacecraft, and Special Sensor Microwave Imager/Sounder (SSMIS) that is flying on the F17 spacecraft, all using the Goddard Profiling Algorithm Version 2010 Version 2 (GPROF2010v2) produced by Colorado State University (see [http://rain.atmos.colostate.edu/RAINMAP10v2/gprof\\_description.html](http://rain.atmos.colostate.edu/RAINMAP10v2/gprof_description.html); Kummerow et al. 2011). Infrared data from the global collection of geosynchronous weather satellites are converted to precipitation estimates using the Precipitation Estimation from Remotely Sensed Information using Artificial Neural Networks Climate Data Record (PERSIANN-CDR), which is produced by NOAA in conjunction with the National Centers for Environmental Information (NCEI; Ashouri et al. 2015). The NASA/GSFC Sounder Research Team provides precipitation estimates from the Television-InfraRed Operational Satellite (TIROS) Operational Vertical Sounder (TOVS) sensors that flew on selected TIROS- and NOAA-series satellites, and on the Advanced Infrared Sounder (AIRS) instrument aboard the Earth Observing System Aqua satellite (Susskind and Pfendtner 1989; Susskind et al. 1997). The TRMM Combined Climatology (TCC) is provided by the University of Maryland. Finally, global gauge analyses are provided by the Deutscher Wetterdienst (DWD) Global Precipitation Climatology Centre (GPCC; Schneider et al. 2017; Schneider et al. 2014; Becker et al. 2013). These five data sets form the basis of the GPCPV3.0 products. These data are merged via a set of algorithms to take advantage of the strengths of each data set and minimize the weaknesses to create a single, best precipitation estimate with associated ancillary fields. A brief algorithm description is provided in the next section.

## 1.2 Algorithm Background

The basis of the GPCPV3.0 algorithm is to use sparse, high-quality GPROF precipitation estimates, which span the range 90°N–90°S (although only considered reliable in the latitude band 60°N–60°S), to calibrate the more frequent, near-global (60°N–60°S) PERSIANN-CDR estimates. GPROF estimates are provided at the orbit level with ~15 orbits/day and the PERSIANN-CDR, based on IR estimates, covers the entire 60°N–60°S area every three hours. The GPROF Level 2 estimates are gridded to the 0.5° resolution, and the PERSIANN-CDR estimates are regridded from the native 0.25° resolution to the 0.5° resolution for consistency.

Calibration is performed by accumulating temporally matched GPROF estimates and PERSIANN-CDR estimates into a multi-level histogram for a calendar month. At the end of the month, a PERSIANN-CDR calibration look-up table is generated based on the histogram-matched

estimates. The 3-hourly PERSIANN-CDR estimates are then calibrated to the GPROF and accumulated for the month. Starting at a seasonally varying boundary around 40°N and S the TOVS/AIRS estimates are increasingly weighted in with the (PMW-calibrated) PERSIANN-CDR, reaching dominant weighting at 58° N and S. Starting at these latitudes (where PERSIANN-CDR becomes less reliable) the TOVS/AIRS estimates are adjusted for use at the high latitudes (58°–90° N and S) to smoothly shift from a zonal average of the combination of PERSIANN-CDR and TOVS/AIRS at 58°N, to a zonal average of the available GPCC values at 70°N. Above 70°N, the zonal-average GPCC adjustment ratio computed at 70°N is used. The corresponding 70°S zonal average is based on a climatological average of a historical collection of GPCC gauges since the gauge population is sparse and sporadic in that zone, and GPCC no longer analyzes precipitation over Antarctica. This field constitutes the satellite-only estimate even though it contains gauge influence at higher latitudes. This field is then adjusted so that it approximately conforms to the (spatially and seasonally varying) TCC, since that is considered the best available long-term average (Wang et al. 2014).

Next, the global satellite-only estimates and the global GPCC gauge analysis are merged. Note that the Legates-Wilmott wind-loss correction is applied to the GPCC gauge analysis prior to merging. The first step in the merge process is to adjust the satellite-only estimate to the large-scale mean of the gauge analysis. The second step is to merge satellite-only estimate with the GPCC gauge analysis based on inverse error variance to produce the final precipitation estimate.

Finally, the field of Probability of Liquid Precipitation Phase is computed based on MERRA2 analyses of surface temperature, humidity, and pressure using a diagnostic lookup table developed by Sims and Liu (2015). Following Sims and Liu, the not-liquid precipitation class is “solid” (snow, graupel, etc.), while the relatively rare “mixed” precipitation class (both liquid and solid in a single observation) is included in “liquid”, since mixed is presumed to generally melt and therefore be a liquid.

Details of the algorithm can be found in Huffman et al. (2019).

## 1.3 Data Disclaimer

This first release, labeled V3.0, is considered to be a beta release, meaning it is useful, but has known limitations. Specifically, the TOVS/AIRS record is not as homogeneous as we expect for a CDR, and the IR input has some noticeable seams. Users should account for these limitations in the use of the V3.0 data. The team continues to work toward improving these issues in a future release.

The GPCPV3.0 data set was carefully constructed to produce global, homogeneous precipitation estimates, so users should use all estimates provided. To determine the relative errors for each grid box, users can examine the satellite-gauge precipitation random error field. Errors over land will vary greatly depending upon the gauge density. For example, areas such as the U.S., Europe, and eastern Australia will have lesser error due to substantial gauge density, whereas



areas such as central Africa, which is deficient in gauge coverage, will have reduced quality. Over ocean, the quality of the input data is fairly uniform, so the associated errors will be correspondingly uniform.

### 1.3.1 Data Citation

The data set source should be acknowledged when the data are used. The American Meteorological Society guidelines contain suggested formats for referencing data sets (<https://www.ametsoc.org/ams/index.cfm/publications/authors/journal-and-bams-authors/formatting-and-manuscript-components/references/dataset-references/>). Following their example for data that are dynamically updated without distinct versions:

Huffman, G.J., D.T. Bolvin, E.J. Nelkin, 2019: GPCP Version 3.0 SG Combined Precipitation Data Set, last updated <date>. GES DISC, Greenbelt, MD, accessed <date>, <https://doi.org/10.5067/TTO0VJF2FSYR>

### 1.3.2 Contact Information

If you have general questions or comments regarding the GPCPV3.0 data set, please contact George Huffman ([george.j.huffman@nasa.gov](mailto:george.j.huffman@nasa.gov)), while questions about accessing and reading the data should be directed to Zhong Liu ([zhong.liu-1@nasa.gov](mailto:zhong.liu-1@nasa.gov)).

## 1.4 What's New?

V3.0 is the inaugural version of the GPCP Version 3 data set. It is considered a beta release, meaning it is useful, but has known limitations. The team continues to work toward improving these issues in a future release.

## 2.0 Data Organization

---

The dataset consists of monthly data, one file per month.

## 2.1 File Naming Convention

**GPCPMON\_L3\_yyyyymm\_V3.0.nc4**

Where:

- yyyy = 4 digit year number [1983 - 2016].
- mm = 2 digit month number [01-12]

Filename example: GPCPMON\_L3\_200702\_V3.0.nc4

## 2.2 File Format and Structure

Data set files are in netCDF version 4 format developed at the University Corporation for Atmospheric Research (UCAR) and Unidata (<https://www.unidata.ucar.edu/software/netcdf/>), with extensions that facilitate the creation of Grid, Point, and Swath data structures. The GPCPV3.0 dataset uses the Grid structure.

## 2.3 Key Science Data Fields

See tables in Section 3.3 for lists of data fields.

# 3.0 Data Contents

---

## 3.1 Dimensions

Each netCDF file contains three dimensions:

latitude – latitude of the center of each row of data (WGS84 reference datum). This one-dimensional array has 360 elements.

longitude – longitude of the center of each column of data (WGS84 reference datum). This one-dimensional array has 720 elements.

time – the begin and end times of the month in minutes since January 1, 1979 00:00. This two-dimensional array has 1 x 2 elements.

## 3.2 Global Attributes

In addition to SDS arrays containing variables and dimension scales, global metadata is

also stored in the files. Some metadata are required by standard conventions, some are provided to meet data provenance requirements, and others are provided as a convenience to users of GPCPV3.0 products. A summary of global attributes present in all files is shown in Table 1.

*Table 1. Global metadata attributes associated with each SDS.*

<b>Attribute Name</b>	<b>Description</b>	<b>Data Type</b>
ShortName	GES DISC short name	string
LongName	GES DISC long name	string
VersionID	version identification number	string
LocalGranuleID	dataset file name	string
Format	dataset format (netCDF-4)	string
RangeBeginningDate	data begin date	string
RangeBeginningTime	data begin time	string
RangeEndingDate	data end date	string
RangeEndingTime	data end time	string
ProductionDateTime	file production date time	string
Conventions	file convention (CF-1.5)	string
Northernmost_Latitude	northernmost latitude contained in the file	float
Southernmost_Latitude	southernmost latitude contained in the file	float
Westernmost_Longitude	westernmost longitude contained in the file	float
Easternmost_Longitude	easternmost longitude contained in the file	float
Latitude_Resolution	latitudinal resolution of the data	float
Longitude_Resolution	longitudinal resolution of the data	float
Entry_ID	GES DISC dataset identifier	string
Entry_Title	GES DISC dataset title	string
Title	description of the dataset	string
Science_Keywords	dataset keywords	string
ISO_Topic_Category	ISO standard dataset category	string
Data_Center_ShortName	GES DISC short name	string
Data_Center_LongName	GES DISC long name	string
Data_Center_URL	GES DISC URL	string
Data_Center_Role	GES DISC role	string
Data_Center_Last_Name	GES DISC support group name	string
Data_Center_Email	GES DISC email	string
Data_Center_Phone	GES DISC phone number	string
Data_Center_Fax	GES DISC Fax number	string
Data_Center_Address	GES DISC address	string
Data_Set_Progress	current state of the dataset	string
Data_Set_Quality	description of the quality of the dataset	string
Summary	description of the dataset	string
Validation_Data	data used to validate the dataset	string
Source	dataset input sources	string
MapProjection	dataset map projection	string
Dataset_Creator	dataset creators	string
Dataset_Title	full dataset title	string
Dataset_Series_Name	dataset name	string
Dataset_Release_Date	release date of current dataset	string
Datset_Release_Place	address of current release	string

Table 1, continued.

Attribute Name	Description	Data Type
Dataset_Publisher	publisher of current dataset	string
IdentifierProductDOI	dataset DOI	string
Data_Presentation_Form	dataset type	string
RelatedURL	alternative URL for the dataset	string
References	dataset references	string
Role	role identifier (science investigator)	string
First_Name	first name of investigator	string
Last_Name	last name of investigator	string
Phone	phone number of science investigator	string
FAX	fax number of science investigator	string
Email	email of science investigator	string
Contact_Address	street address of science investigator	string
City	city of science investigator	string
Province_or_State	state of science investigator	string
Postal_Code	zip code of science investigator	string
Country	country of science investigator	string
Use_Constraints	constraints on the use of the dataset	string
Distribution_Media	form of dataset access	string
Distribution_Size	size of dataset file	string
Distribution_Format	dataset format (netCDF-4)	string
IdentifierProductDOIAuthority	DOI authority	string
ProcessingLevel	dataset processing level	string
Institution	where the data were produced	string
Fees	costs for use of the dataset	string

A list of key metadata fields is given in Table 2. These and other metadata fields can be found in the "**CoreMetadata.0**" global attribute. Global attributes in a GPCPV3.0 file can be viewed with *ncdump* software as described in Section 4.1 [*ncdump -h -c <GPCPV3.0 file>*].

Table 2. Key Metadata Items

Name	Type	Description
_FillValue	float or short int	Value used to identify missing data.
long_name	string	ad hoc description of the variable.
valid_range	float or short int	valid data value ranges for the variable
Units	string	The units of the variable.
cell_methods	String	Temporal unit of data

## 3.3 Products/Parameters

The list of data fields contained in each GPCP V3.0 Monthly dataset is given in Table 3, together with brief descriptions and native units.

Table 3. Data fields in the GPCP V3.0 Monthly dataset. The latitude/longitude use the WGS84 reference datum.

Data Field Name	Description	Units
latitude	latitude of the center of the grid element	degrees_north
longitude	longitude of the center of the grid element	degrees_east
time	begin date and time of the data	minutes since 1979-01-01 00:00:00
sat_gauge_precip	combined satellite-gauge precipitation estimate	mm/day
sat_gauge_error	combined satellite-gauge precipitation random error estimate	mm/day
satellite_precip	satellite-only precipitation estimate	mm/day
satellite source	source of the contributing satellite estimate *	index values
gauge_precip	wind-loss adjusted gauge precipitation	mm/day
probability_liquid_precip	probability of liquid phase precipitation	percent

\* The satellite source field has the index values 0 = PMW-adj. IR, 2 = blend of PMW-adj. IR and TOVS/AIRS, 4 = TOVS/AIRS.

## 4.0 Options for Reading the Data

---

It is important that the GPCPV3.0 is as accessible as possible to a wide range of users. The MEASUREs project selected netCDF as it is designed to store scientific data and is easily accessible with command-line tools and standard application packages. If you have trouble reading the data please contact Zhong Liu at [zhong.liu-1@nasa.gov](mailto:zhong.liu-1@nasa.gov).

### 4.1 Command Line Utilities

There are several built-in netCDF utilities that are available to access the GPCPV3.0 data. See <https://www.unidata.ucar.edu/software/netcdf> for a complete list of netCDF command-line utilities. The most useful is *ncdump*:

#### ***ncdump***

The *ncdump* tool can be used as a simple browser for netCDF data files, to display the dimension names and sizes; variable names, types, and shapes; attribute names and values. The most common use of *ncdump* is with the `-h` option, in which only the header information is displayed.

ncdump [-c|-h] [-v ...] [[-b|-f] [c|f]] [-l len] [-n name] [-d n[,n]] filename

Options/Arguments:

- [-c] Coordinate variable data and header information
  - [-h] Header information only, no data
  - [-v var1[,...]] Data for variable(s) <var1>,... only data
  - [-f [c|f]] Full annotations for C or Fortran indices in data
  - [-l len] Line length maximum in data section (default 80)
  - [-n name] Name for netCDF (default derived from file name)
  - [-d n[,n]] Approximate floating-point values with less precision
- filename File name of input netCDF file

## 4.2 Tools/Programming

Any application package that supports the netCDF file format can be used to read these data. If you are new to netCDF or to the GPCP V3.0 datasets, the following tool might be a useful starting point:

- Panoply (<https://www.giss.nasa.gov/tools/panoply/>)

For more advanced users or programmers:

- IDL (<http://www.harrisgeospatial.com/SoftwareTechnology/IDL.aspx>)
- Matlab (<http://www.mathworks.com/products/matlab/>)
- GrADS (<http://cola.gmu.edu/grads/>)
- Python
- Java
- C/C++
- Fortran77/90

As well, the GPCP V3.0 data are accessible through the Giovanni web-based visualization and analysis application, <https://giovanni.gsfc.nasa.gov/giovanni/>.

## 5.0 Data Services

---

The data are currently available through the **Data Holdings** page of the **MEaSUREs portal** at <https://disc.gsfc.nasa.gov/measures>.

If you need assistance or wish to report a problem:

**Email:** [gsfc-dl-help-disc@mail.nasa.gov](mailto:gsfc-dl-help-disc@mail.nasa.gov)

**Voice:** 301-614-5224

**Fax:** 301-614-5268

**Address:**

Goddard Earth Sciences Data and Information Services Center NASA Goddard Space Flight Center Code 610.2 Greenbelt, MD 20771 USA

## 6.0 More Information

---

For more information on our project or data, please refer to the Algorithm Theoretical Basis Document (Huffman et al. 2019).

## 7.0 Acknowledgements

---

This project is funded by NASA's Making Earth Science Data Records for Use in Research Environments (MEaSUREs) Program and managed by Dr. Lucia Tsaoussi.

## References

Ashouri, H., K-L. Hsu, S. Sorooshian, D.K. Braithwaite, K.R. Knapp, L.D. Cecil, B.R. Nelson, O.P. Prat, 2015: PERSIANN-CDR, Daily Precipitation Climate Data Record from Multisatellite Observations for Hydrological and Climate Studies. *Bull. Amer. Meteor. Soc.*, 96, 69-83. [doi:10.1175/BAMS-D-13-00068.1](https://doi.org/10.1175/BAMS-D-13-00068.1)

Becker, A., P. Finger, A. Meyer-Christoffer, B. Rudolf, K. Schamm, U. Schneider, M. Ziese, 2013: A Description of the Global Land-Surface Precipitation Data Products of the Global Precipitation Climatology Centre with Sample Applications Including Centennial (Trend) Analysis from 1901–Present. *Earth Syst. Sci. Data*, 5, 71-99. [doi:10.5194/essd-5-71-2013](https://doi.org/10.5194/essd-5-71-2013)

Huffman, G.J., R.F. Adler, D.T. Bolvin, K. Hsu, C. Kidd, E.J. Nelkin, J. Tan, P. Xie, 2019: Algorithm Theoretical Basis Document (ATBD) for Global Precipitation Climatology Project Version 3.0 Precipitation Data. MEaSUREs project, Greenbelt, MD, 27 pp. Available online: [https://docserver.gesdisc.eosdis.nasa.gov/public/project/MEaSUREs/GPCP/GPCP\\_ATBD\\_V3.0.pdf](https://docserver.gesdisc.eosdis.nasa.gov/public/project/MEaSUREs/GPCP/GPCP_ATBD_V3.0.pdf) (last accessed: March 14, 2019).

- Kummerow, C.D., S. Ringerud, J. Crook, D. Randel, W. Berg, 2011: An Observationally Generated A Priori Database for Microwave Rainfall Retrievals. *J. Atmos. Oc. Technol.*, **28**, 113-130. doi:10.1175/2010JTECHA1468.1
- Schneider, U., A. Becker, P. Finger, A. Meyer-Christoffer, M. Ziese, B. Rudolf, 2014: GPCC's New Land Surface Precipitation Climatology based on Quality-controlled In Situ Data and its Role in Quantifying the Global Water Cycle. *Theor. Appl. Climatol.*, **115**, 15, doi:10.1007/s00704-013-0860-x
- Schneider, U., P. Finger, A. Meyer-Christoffer, E. Rustemeier, M. Ziese, A. Becker, 2017: Evaluating the Hydrological Cycle over Land Using the Newly-Corrected Precipitation Climatology from the Global Precipitation Climatology Centre (GPCC). *Atmosphere*, **8**, 52, doi:10.3390/atmos8030052
- Sims, E.M., and G. Liu, 2015: A Parameterization of the Probability of Snow–Rain Transition. *J. Hydrometeor.*, **16**, 1466–1477. doi:10.1175/JHM-D-14-0211.1
- Susskind, J. C.D. Barnet, J.M. Blaisdell, 2003: Retrieval of Atmospheric and Surface Parameters from AIRS/AMSU/HSB Data in the Presence of Clouds. *IEEE Trans. Geosci. Rem. Sens.*, **41**, 390-409. doi:10.1109/TGRS.2002.808236
- Susskind, J.M. Blaisdell, L. Iredell, 2014: Improved Methodology for Surface and Atmospheric Soundings, Error Estimates, and Quality Control Procedures: The Atmospheric Infrared Sounder Science Team Version-6 Retrieval Algorithm. *J. Appl. Rem. Sens.*, **8**, Art. No. 084994. doi:10.1117/1.JRS.8.084994
- Susskind, J., J. Pfaendtner, 1989: Impact of Interactive Physical Retrievals on NWP. *Report on the Joint ECMWF/EUMETSAT Workshop on the Use of Satellite Data in Operational Weather Prediction: 1989-1993, Vol. 1*, T. Hollingsworth, Ed., ECMWF, Shinfield Park, Reading RG2 9AV, U.K., 245-270.
- Susskind, J., P. Piraino, L. Rokke, L. Iredell, A. Mehta, 1997: Characteristics of the TOVS Pathfinder Path A Dataset. *Bull. Amer. Meteor. Soc.*, **78**, 1449-1472.
- Wang, J., R.F. Adler, G.J. Huffman, and D. Bolvin, 2014: An Updated TRMM Composite Climatology of Tropical Rainfall and Its Validation. *J. Climate*, **27**, 273–284. doi:10.1175/JCLI-D-13-00331.1
- Wu, H., R.F. Adler, Y. Tian, G.J. Huffman, H. Li, J. Wang, 2014: Real-Time Global Flood Estimation Using Satellite-Based Precipitation and a Coupled Land Surface and Routing Model, *Water Resour. Res.*, **50**, 2693–2717. doi:10.1002/2013WR014710